ABclonal® www.abclonal.com

Phospho-c-Jun-S63 Rabbit pAb

Catalog No.: AP1190 1 Publications

Basic Information

Observed MW

44kDa

Calculated MW

36kDa

Category

Polyclonal Antibody

Applications

WB,IHC-P,ELISA

Cross-Reactivity

Human, Mouse, Rat

Background

This gene is the putative transforming gene of avian sarcoma virus 17. It encodes a protein which is highly similar to the viral protein, and which interacts directly with specific target DNA sequences to regulate gene expression. This gene is intronless and is mapped to 1p32-p31, a chromosomal region involved in both translocations and deletions in human malignancies.

Recommended Dilutions

WB 1:500 - 1:1000

IHC-P 1:50 - 1:200

ELISA Recommended starting

concentration is 1 µg/mL. Please optimize the concentration based on your specific

assay requirements.

Immunogen Information

Gene IDSwiss Prot
P05412

Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

Synonyms

AP1; p39; AP-1; cJUN; c-Jun; Phospho-c-Jun-S63

Contact

www.abclonal.com


Product Information

SourceIsotypePurificationRabbitIgGAffinity purification

Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.



Western blot analysis of lysates from 293T cells, using Phospho-c-Jun-S63 Rabbit pAb (AP1190) at 1:1000 dilution. 293T cells were treated with Anisomycin (25 μ g/mL) at 37°C for 30 minutes after serum-starvation overnight.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.


Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).


Exposure time: 30s.



Immunohistochemistry analysis of paraffin-embedded Rat brain using Phospho-c-Jun-S63 Rabbit pAb (AP1190) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Human breast cancer using Phospho-c-Jun-S63 Rabbit pAb (AP1190) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.



Immunohistochemistry analysis of paraffin-embedded Mouse brain using Phospho-c-Jun-S63 Rabbit pAb (AP1190) at dilution of 1:200 (40x lens). High pressure antigen retrieval performed with 0.01M Citrate buffer (pH 6.0) prior to IHC staining.