

AP1504

Leader in Biomolecular Solutions for Life Science



# Phospho-p53-S15 Rabbit mAb

Catalog No.: AP1504

Recombinant

## Basic Information

### Observed MW

53kDa

### Calculated MW

44kDa

### Category

SMab Recombinant Monoclonal  
Antibody

### Applications

WB,IF/ICC,ELISA

### Cross-Reactivity

Human

### CloneNo number

ARC63388

## Recommended Dilutions

**WB** 1:1000 - 1:3000

**IF/ICC** 1:50 - 1:200

**ELISA** Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.

## Contact

 [www.abclonal.com](http://www.abclonal.com)

## Background

This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and oligomerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancers such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters result in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from the use of alternate translation initiation codons from identical transcript variants (PMIDs: 12032546, 20937277).

## Immunogen Information

### Gene ID

7157

### Swiss Prot

P04637

### Immunogen

Synthetic peptide. This information is considered to be commercially sensitive.

### Synonyms

P53; BCC7; LFS1; BMFS5; TRP53; Phospho-p53-S15

## Product Information

### Source

Rabbit

### Isotype

IgG

### Purification


Affinity purification

### Storage

Store at -20°C. Avoid freeze / thaw cycles.

Buffer: PBS containing 50% glycerol and 0.05% BSA, preserved with proclin300 or sodium azide (as specified on the Certificate of Analysis), pH 7.3.

## Validation Data



Western blot analysis of lysates from 293T cells, using Phospho-p53-S15 Rabbit mAb (AP1504) at 1:3000 dilution. 293T cells were treated with UV at room temperature for 15-30 minutes.


Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (AS014) at 1:10000 dilution.

Lysates/proteins: 25µg per lane.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit (RM00020).

Exposure time: 20s.



Confocal imaging of 293T treated with UV and 293T cells using Phospho-p53-S15 Rabbit mAb (AP1504) at dilution of 1:200 (1000x lens). Blue: DAPI for nuclear staining.