

Recombinant Human IL-4R alpha/CD124 Protein

Catalog No.: RP00099 Recombinant

Sequence Information

Species Gene ID Swiss Prot HEK293 cells 3566 P24394

Tags

C-His

Synonyms

IL4R;CD124;IL-4RA;IL4RA

Product Information

Source Purification
HEK293 cells > 87% by SDSPAGE.

Endotoxin

 $< 0.1 \; \text{EU/}\mu\text{g}$ of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

www.abclonal.com

Background

This protein is alpha chain of the interleukin-4 receptor, a type I transmembrane protein that can bind interleukin 4 and interleukin 13 to regulate IgE production. The encoded protein also can bind interleukin 4 to promote differentiation of Th2 cells. A soluble form of the encoded protein can be produced by proteolysis of the membrane-bound protein, and this soluble form can inhibit IL4-mediated cell proliferation and IL5 upregulation by T-cells.

Basic Information

Description

Recombinant Human IL-4R alpha/CD124 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gly24-His232) of human IL4R/CD124 (Accession #NP_000409.1) fused with a 6×His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized recombinant human IL4R at 2 μ g/mL (100 μ L/well) can bind recombinant human IL4. The ED₅₀ of recombinant human IL4R is 2-4 ng/mL.


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

-20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Immobilized recombinant human IL4R at 2 μ g/mL (100 μ L/well) can bind recombinant human IL4. The ED₅₀ of recombinant human IL4R is 2-4 ng/mL.