ABclonal www.abclonal.com

Recombinant Human Basigin/EMMPRIN/CD147 Protein

Catalog No.: RP00104 Recombinant

Sequence Information

Species Gene ID Swiss Prot HEK293 cells 682 P35613

Tags

C-hFc&His

Synonyms

5F7; CD147; EMMPRIN; OK; TCSF;BSG;

basigin;CD147;EMMPRIN;OK;TCSF;basigin

Product Information

Source Purification
HEK293 cells > 95% by SDSPAGE.

Endotoxin

 $< 0.1 \; \text{EU/}\mu\text{g}$ of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

This protein is a plasma membrane protein that is important in spermatogenesis, embryo implantation, neural network formation, and tumor progression. The encoded protein is also a member of the immunoglobulin superfamily. Multiple transcript variants encoding different isoforms have been found for this gene.

Basic Information

Description

Recombinant Human Basigin/EMMPRIN/CD147 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Thr25-His205) of human CD147/EMMPRIN/Basigin (Accession #NP_001719.2) fused with an Fc, 6×His tag at the C-terminus.

Bio-Activity


Measured by its binding ability in a functional ELISA.Immobilized Human CD147 at 1μ g/mL (100 μ L/well) can bind CD147 Mouse mAb with a linear range of 0.12-3.05 ng/ml.

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.
 -80°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human Basigin/CD147 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 65-75 kDa.



Immobilized Human CD147 at $1\mu g/mL$ (100 $\mu L/well$) can bind CD147 Mouse mAb with a linear range of 0.12-3.05 ng/ml.