

Recombinant Human BTN1A1/Butyrophilin Protein

Catalog No.: RP00249 Recombinant

Sequence Information

Species Gene ID Swiss Prot HEK293 cells 696 Q13410

Tags

C-His

Synonyms

BTN1A1;BT;BTN;BTN1

Product Information

Source Purification

HEK293 cells ≥ 95 % as determined by SDS-PAGE.

Endotoxin

 $< 0.1 \; \text{EU/}\mu\text{g}$ of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

www.abclonal.com

Background

Basic Information

Description

Recombinant Human BTN1A1/Butyrophilin Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala 27 - Arg 242) of human BTN1A1/Butyrophilin (Accession $\#NP_001723.2$) fused with a 6×His tag at the C-terminus.

Bio-Activity

Measured by its ability to inhibit Anti-CD3-induced proliferation of jurkat cells. The ED_{50} for this effect is 3.35-13.4 ng/mL.

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.


Avoid repeated freeze/thaw cycles.

^{*} For your safety and health, please wear a lab coat and disposable gloves when handling.


Validation Data



Recombinant Human BTN1A1/Butyrophilin Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Recombinant Human BTN1A1 inhibits Anti-CD3-induced proliferation of jurkat cells. The ED_{so} for this effect is 6.6-26.4ng/mL.



Recombinant Human BTN1A1 inhibits Anti-CD3-induced proliferation of jurkat cells. The ED_{so} for this effect is 3.35-13.4ng/mL.