

RP00270

Leader in Biomolecular Solutions for Life Science



# Recombinant Human CSF3R/G-CSF-R/CD114 Protein

Catalog No.: RP00270

Recombinant

## Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	1441	Q99062

### Tags

C-His

### Synonyms

CSF3R;CD114;GCSFR;SCN7

## Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

### Endotoxin

< 0.1 EU/μg of the protein by LAL method.

### Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

## Background

### Basic Information

#### Description

Recombinant Human CSF3R/G-CSF-R/CD114 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Glu25-Pro621) of human GCSF Receptor (Accession #NP\_000751.1) fused with a 6×His tag at the C-terminus.

#### Bio-Activity

Measured by its ability to inhibit the GCSF-induced proliferation of M-NFS-60 mouse myelogenous leukemia lymphoblast cells. The ED<sub>50</sub> for this effect is typically 31.5-126 ng/mL.

#### Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

## Contact




[www.abclonal.com](http://www.abclonal.com)

Validation Data



Recombinant Human CSF3R/G-CSF-R/CD114 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 90-105 kDa.



Recombinant Human GCSF Receptor inhibits the GCSF-induced proliferation of M-NFS-60 mouse myelogenous leukemia lymphoblast cells. The ED<sub>50</sub> for this effect is typically 31.5-126 ng/mL.