ABclonal www.abclonal.com

Recombinant Human Trk-C/NTRK3 Protein

Catalog No.: RP00288 Recombinant

Sequence Information

Species Gene ID Swiss Prot HEK293 cells 4916 Q16288

Tags

C-hFc&His

Synonyms

NTRK3;GP145-TrkC;TRKC;gp145(trkC)

Product Information

Source Purification HEK293 cells > 97% by SDS-PAGE.

Endotoxin

< 1.0 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

 \odot

www.abclonal.com

Background

NT-3 growth factor receptor also known as neurotrophic tyrosine kinase receptor type 3 or TrkC tyrosine kinase or Trk-C receptor, is a member of the neurotrophic tyrosine receptor kinase (NTRK) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. Signalling through this kinase leads to cell differentiation and may play a role in the development of proprioceptive neurons that sense body position. Mutations in TrkC encoding gene have been associated with medulloblastomas, secretory breast carcinomas and other cancers.

Basic Information

Description

Recombinant Human Trk-C/NTRK3 Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Cys32-Asp428) of human TrkC/NTRK3 (Accession #NP_001012338.1) fused with an Fc, 6×His tag at the Cterminus.

Bio-Activity


Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human NTF3 at 2 μ g/mL (100 μ L/well) can bind Recombinant human TrkC with a linear range of 22-90 ng/mL.

Storage


Store at -20° C. Store the lyophilized protein at -20° C to -80° C up to 1 year from the date of receipt.

 After reconstitution, the protein solution is stable at -20° C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Human Trk-C/NTRK3 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 120 kDa.



Immobilized Recombinant Human NTF3 at 2 μ g/mL (100 μ L/well) can bind Recombinant human TrkC with a linear range of 22-90 ng/mL.