

RP00385

Leader in Biomolecular Solutions for Life Science



Recombinant Human ICOSLG/B7-H2/CD275 Protein

Catalog No.: RP00385

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	23308	O75144

Tags

N-hFC

Synonyms

ICOSLG;B7-H2;B7H2;B7RP-1;B7RP1;CD275;GL50;ICOS-L;ICOSL;LICOS

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Endotoxin

< 0.01 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.2 μm filtered solution of PBS,pH7.4.

Reconstitution

Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water.

Background

Inducible co-stimulator ligand (ICOSL) is a member of the B7 family of co-stimulatory molecules related to B7-1 and B7-2. It is a transmembrane glycoprotein with extracellular IgV and IgC domains and binds to ICOS on activated T cells, thus delivers a positive costimulatory signal for optimal T cell function. The structural features of ICOSL are crucial for its costimulatory function. The present study shows that ICOSL displays a marked oligomerization potential, resembling more like B7-1 than B7-2. B7-H2-dependent signaling may play an active role in a proliferative response rather than in cytokine and chemokine production. The CD28/B7 and ICOS/B7-H2 pathways are both critical for costimulating T cell immune responses. Deficiency in either pathway results in defective T cell activation, cytokine production, and germinal center formation.

Basic Information

Description

Recombinant Human ICOSLG/B7-H2/CD275 Protein is produced by Human Cell expression system. The target protein is expressed with sequence (Asp19-Ser258) of human ICOSLG/B7-H2/CD275 (Accession #O75144) fused with a 6×His tag at the C-terminus.

Bio-Activity

Storage

Store the lyophilized protein at -20°C to -80 °C for long term.
After reconstitution, the protein solution is stable at -20 °C for 3 months, at 2-8 °C for up to 1 week.
Avoid repeated freeze/thaw cycles.

Contact



www.abclonal.com