

RP00422

Leader in Biomolecular Solutions for Life Science



Recombinant Human TROP-2/TACSTD2 Protein

Catalog No.: RP00422

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	4070	P09758

Tags

C-hFc

Synonyms

EGP1; EGP-1; TROP2; GA733-1; gp50; T16; TACSTD2; TROP-2; M1S1; TACD2

Product Information

Source

HEK293 cells

Purification

> 95% by SDS-PAGE ; > 95% by HPLC

Endotoxin

< 1 EU/μg of the protein by LAL method

Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Trop-2, also known as epithelial glycoprotein-1 antigen (EGP-1), is a protein that in humans is encoded by the TACSTD2 gene. Mutations of this gene result in gelatinous drop-like corneal dystrophy, an autosomal recessive disorder characterized by severe corneal amyloidosis leading to blindness.

Basic Information

Description

Recombinant Human TROP-2/TACSTD2 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (His27-Thr274) of Human TROP-2/TACSTD2 (Accession #P09758) fused with a C-hFc tag at the C-terminus.

Bio-Activity

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

Contact




www.abclonal.com


Validation Data



Human TROP-2 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



The purity of Human TROP-2 is greater than 95% as determined by SEC-HPLC.



Immobilized Human TROP-2, hFc Tag at 0.5 μg/mL (100 μL/well) on the plate. Dose response curve for Biotinylated Anti-TROP-2 Antibody, hFc Tag with the EC₅₀ of 11.3ng/mL determined by ELISA.