

RP00527

Leader in Biomolecular Solutions for Life Science



Recombinant Human TNFSF7/CD27 Ligand/CD70 Protein

Catalog No.: RP00527 **Recombinant**

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	970	P32970-1

Tags

N-hFc

Synonyms

CD70 molecule; CD70; TNFSF7; CD27 Ligand; CD27-L; CD27LG; ?Ki-24 antigen; TNFSF7G

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE ; > 95% by HPLC

Endotoxin

< 1 EU/μg of the protein by LAL method

Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

CD70, also named CD27 ligand (CD27L), is a type II transmembrane glycoprotein belonging to the TNF superfamily (TNFSF) and has been designated TNFSF7. CD70 is a cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells.

Basic Information

Description

Recombinant Human CD27 Ligand/CD70 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Gln39-Pro193) of Human TNFSF7/CD27 Ligand/CD70 (Accession #P32970-1) fused with tag at the N-terminus.

Bio-Activity

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

Contact




www.abclonal.com


Validation Data



Recombinant Human TNFSF7/CD27 Ligand/CD70 Protein was determined by Tris-Bis PAGE under reducing conditions.



The purity of Human CD27 Ligand is greater than 95% as determined by SEC-HPLC.



Immobilized Human CD27, His Tag at 0.5 μg/mL (100 μL/well) on the plate. Dose response curve for Human CD27 Ligand, hFc Tag with the EC₅₀ of 14.2ng/mL determined by ELISA.