# ABclonal www.abclonal.com

## **Recombinant Human BST-1/CD157 Protein**

Catalog No.: RP00541 Recombinant

#### **Sequence Information**

**Species Gene ID Swiss Prot** HEK293 cells 683 Q10588

**Tags** C-His

Synonyms BST1;CD157

#### **Product Information**

**Source** Purification HEK293 cells > 97% by SDS-PAGE.

#### **Endotoxin**

< 0.01 EU/µg of the protein by LAL method.

#### **Formulation**

Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.Contact us for customized product form or formulation.

#### Reconstitution

Centrifuge the tube before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

#### **Contact**

€

www.abclonal.com

#### **Background**

The cluster of differentiation (CD) system is a glycosyl phosphatidylinositol anchored membrane protein thatbelongs to the CD38 family.It is generally used in immunophynotyping. CD157 was discovered in a bonemarrow stromal cell line where it facilitates pre-B-cell growth. CD157 is a bifunctional ectoenzyme that exhibitsboth ADP-ribosyl cyclase and cyclic ADP ribose hydrolase activities followed with CD38. It plays a role inrheumatoid arthritis (RA) due to its enhanced expression in RA-derived bone marrow stromal cell lines. Studieshave shown that this protein have a role in predicted to function as a cell surface receptor and animmunoregulatory molecule.

#### **Basic Information**

#### **Description**

Recombinant Human BST-1/CD157 Protein is produced by Human Cells expression system. The target protein is expressed with sequence (Gly29-Lys292) of human CD157/BST1/ADP-ribosyl cyclase 2/Cyclic ADP-ribose hydrolase 2 (Accession #Q10588) fused with a 6×His tag at the C-terminus.


#### **Bio-Activity**

#### Storage

Store at  $-20^{\circ}$ C. Store the lyophilized protein at  $-20^{\circ}$ C to  $-80^{\circ}$ C up to 1 year from the date of receipt. <br/>
-20^{\circ} for 3 months, at 2-8^{\circ} for up to 1 week.

Avoid repeated freeze/thaw cycles.

### **Validation Data**



Recombinant Human BST-1/CD157 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 35-45 kDa.