

RP00601

Leader in Biomolecular Solutions for Life Science



Recombinant Human CD3 epsilon Protein

Catalog No.: RP00601

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	916	P07766

Tags

C-hFc

Synonyms

CD3e; CD3E; T3E; FLJ18683; TCRE;
CD3-epsilon; IMD18

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE

Endotoxin

< 1 EU/μg of the protein by LAL method

Formulation

Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

T-cell surface glycoprotein CD3 epsilon & CD3 gamma chain, also known as CD3E & CD3G, are single-pass type I membrane proteins. When antigen presenting cells (APCs) activate T-cell receptor (TCR), TCR-mediated signals are transmitted across the cell membrane by the CD3 chains CD3D, CD3E, CD3G and CD3Z. All CD3 chains contain immunoreceptor tyrosine-based activation motifs (ITAMs) in their cytoplasmic domain.

Basic Information

Description

Recombinant Human CD3 epsilon Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Asp23-Asp126) of Human CD3 epsilon (Accession #P07766) fused with a C-hFc tag at the C-terminus.

Bio-Activity


Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


Contact



www.abclonal.com



Human CD3E on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.



Immobilized Human CD3E, hFc Tag at 1 µg/mL (100 µL/Well) on the plate. Dose response curve for Biotinylated Anti-CD3 Antibody, hFc Tag with the EC₅₀ of 33.5ng/mL determined by ELISA (QC Test).