ABclonal www.abclonal.com

Recombinant Human CTLA-4/CD152 Protein

Catalog No.: RP01044 Recombinant

Sequence Information

Species Gene ID Swiss Prot HEK293 cells 1493 P16410

Tags

C-His

Synonyms

CTLA4;ALPS5;CD;CD152;CELIAC3;CTLA-4;GRD4;GSE;IDDM12

Product Information

Source Purification HEK293 cells > 90% by SDS-

PAGE.

Endotoxin

< 1.0 EU/ μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

€

www.abclonal.com

Background

Basic Information

Description

Recombinant Human CTLA-4/CD152 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala37-Phe162) of human CTLA-4 (Accession #NP_005205.2) fused with a 6×His tag at the C-terminus

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Ipilimumab at 1 μ g/mL (100 μ L/well) can bind Recombinant Human CTLA-4 with a linear range of 6-18 ng/mL.|2.Measured by its binding ability in a functional ELISA.Immobilized PE anti-human CD152 (CTLA-4) Antibody at 1 μ g/mL (25 μ L/well) can bind Human CTLA-4 with a linear range of 0.46-2.05 ng/mL.


Storage

Store at -20° C. Store the lyophilized protein at -20° C to -80° C up to 1 year from the date of receipt.


-20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.


Validation Data




Recombinant Human CTLA-4/CD152 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 25 kDa.




The purity of Human CTLA-4 Protein (Cat.RP01044) was greater than 90% as determined by SEC-HPLC.



Immobilized Recombinant Mouse CD86 at 2 μ g/mL (100 μ L/well) can bind Recombinant Human CTLA-4 with a linear range of 15-50 ng/mL.



Immobilized Ipilimumab at 1 μ g/mL (100 μ L/well) can bind Recombinant Human CTLA-4 with a linear range of 6-18 η g/mL.



Immobilized PE anti-human CD152 (CTLA-4) Antibody at 1 μ g/mL (25 μ L/well) can bind Human CTLA-4 with a linear range of 0.46-2.05 ng/mL.