

Recombinant Mouse c-Kit ligand/KITLG/SCF Protein

Catalog No.: RP01055 Recombinant 1 Publications

Sequence Information

Species Gene ID Swiss Prot HEK293 cells 17311 P20826-1

Tags

C-His

Synonyms

KITLG;FPH2;KL-1;Kitl;MGF;SCF;SF;SHEP 7;KL;SCF/KITLG

Product Information

Purification Source HEK293 cells > 95% by SDS-

PAGE.

Endotoxin

< 0.1 EU/µg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

•

www.abclonal.com

Background

Basic Information

Description

Recombinant Mouse c-Kit ligand/KITLG/SCF Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Lys26-Ala189) of mouse SCF (Accession #NP_038626.1.) fused with a 6×His tag at the Cterminus.

Bio-Activity


1.Measured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED₅₀ for this effect is 0.5-1.0 ng/mL, corresponding to a specific activity of 1.0×10⁶-2.0×10⁶ units/mg.|2. Measured by its binding ability in a functional ELISA.Immobilized recombinant Mouse SCF at 2 µg/mL (100 µL/well) can bind recombinant human CD117 with a linear range of 1.5-10 ng/mL.

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.
 After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant Mouse c-Kit ligand/KITLG/SCF Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Immobilized recombinant Mouse SCF at 2 μ g/mL (100 μ L/well) can bind recombinant human CD117 with a linear range of 1.5-10ng/mL.