

# Recombinant Mouse Lipocalin-2/NGAL/LCN2 Protein

Catalog No.: RP01064 Recombinant

# **Sequence Information**

**Species Gene ID Swiss Prot** HEK293 cells 16819 P11672

**Tags** 

C-His

**Synonyms** 

NGAL;LCN2;Lipocalin-2;24p3;MSFI;LCN 2

### **Product Information**

**Source** Purification
HEK293 cells > 95% by SDS-

PAGE.

### **Endotoxin**

 $< 0.1 \; \text{EU/}\mu\text{g}$  of the protein by LAL method.

### **Formulation**

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.Contact us for customized product form or formulation.

#### Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

### **Contact**

 $oldsymbol{\Theta}$ 

www.abclonal.com

# **Background**

## **Basic Information**

### Description

Recombinant Mouse Lipocalin-2/NGAL Protein is produced by HEK293 expression system. The target protein is expressed with sequence (Gln21-Asn200) of mouse Lipocalin-2/NGAL (Accession #NP\_032517.1) fused with an 8×His tag at the C-terminus.

### **Bio-Activity**


The binding of Fe(DHBA)3 results in the quenching of Trp fluorescence in recombinant mouse Lipocalin-2. Recombinant mouse Lipocalin-2 can bind >1.65  $\mu$ M of Fe(DHBA)3.

### Storage


Store at  $-20^{\circ}$ C. Store the lyophilized protein at  $-20^{\circ}$ C to  $-80^{\circ}$ C up to 1 year from the date of receipt. <br/>
-20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

# **Validation Data**



Recombinant Mouse Lipocalin-2/NGAL/LCN2 Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



The binding of Fe(DHBA)3 results in the quenching of Trp fluorescence in recombinant mouse Lipocalin-2. Recombinant mouse Lipocalin-2 can bind >1.65  $\mu$ M of Fe(DHBA)3.