

RP01160

Leader in Biomolecular Solutions for Life Science



Recombinant Human IL-23/IL-12B&IL-23A Protein

Catalog No.: RP01160

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	3593&51561P29460&Q9	NPF7

Tags

No tag(IL12B)&N-his(IL23A)

Synonyms

IL-23 p19/IL-12 p40;IL23;SGRF ;IL-23
alpha & IL-12 beta

Product Information

Source	Purification
HEK293 cells	> 95% by SDS- PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL
method.

Formulation

Lyophilized from a 0.22 μm filtered
solution of PBS, pH 7.4.Contact us for
customized product form or
formulation.

Reconstitution

Centrifuge the vial before opening.
Reconstitute to a concentration of
0.1-0.5 mg/mL in sterile distilled water.
Avoid vortex or vigorously pipetting
the protein. For long term storage, it is
recommended to add a carrier protein
or stabilizer (e.g. 0.1% BSA, 5% HSA,
10% FBS or 5% Trehalose), and aliquot
the reconstituted protein solution to
minimize free-thaw cycles.

Background

Basic Information

Description

Active Recombinant Human IL-12B&IL-23A Protein is produced by HEK293 cells
expression system. The target protein is expressed with sequence (Ile23-
Ser328&Arg20-Pro189) of human IL-23 (Accession #NP_002178.2/NP_057668.1)
fused with a 6×His tag at the N-terminus(IL23A).

Bio-Activity

1.Measured by its binding ability in a functional ELISA.Immobilized Human
IL-12B&IL-23A at 5μg/mL (100 μL/well) can bind Human IL23R with a linear range
of 4-135 ng/mL.2.Measured by its ability to induce IL-17 secretion by mouse
splenocytes. The ED₅₀ for this effect is 21.23-84.94 ng/mL,
corresponding to a specific activity of
1.17×10⁴~4.71×10⁴ units/mg.

Storage


Store at -20°C.Store the lyophilized protein at -20°C to -80 °C up to 1 year from
the date of receipt. After reconstitution, the protein solution is stable at
-20°C for 3 months, at 2-8°C for up to 1 week.
Avoid repeated freeze/thaw cycles.

Contact




www.abclonal.com


Validation Data



Recombinant Human IL-23/IL-12B&IL-23A Protein was determined by SDS-PAGE under reducing (R) and non-reducing (NR) conditions.



Immobilized Human IL-12B&IL-23A at 5µg/mL (100 µL/well) can bind Human IL23R with a linear range of 4-135 ng/mL.



Recombinant Human IL-23/IL-12B&IL-23A induce IL-17 secretion by mouse splenocytes. The ED₅₀ for this effect is 21.23-84.94 ng/mL, corresponding to a specific activity of 1.17×10⁴~4.71×10⁴ units/mg.