

RP01218

Leader in Biomolecular Solutions for Life Science

Recombinant Human TNFRSF5/CD40 Protein

Catalog No.: RP01218

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	958	P25942-1

Tags

C-hFc&His

Synonyms

CD40;Bp50;CDW40;TNFRSF5;p50;
Bp50; CDW40; TNFRSF5

Product Information

Source	Purification
HEK293 cells	> 97% by SDS-PAGE.

Endotoxin

< 0.1 EU/μg of the protein by LAL method.

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4. Contact us for customized product form or formulation.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Basic Information

Description

Recombinant Human TNFRSF5/CD40 Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Glu21-Arg193) of human CD40/TNFRSF5 (Accession #NP_001241.1) fused with a Fc, 6×His tag at the C-terminus.

Bio-Activity

Measured by its binding ability in a functional ELISA. Immobilized Recombinant Human CD40 Ligand at 2 μg/mL (100 μL/well) can bind Recombinant Human CD40 with a linear range of 224-898 ng/mL.

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

Contact




www.abclonal.com


Validation Data



Recombinant Human TNFRSF5/CD40 Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 55-60 kDa.



The purity of human CD40/TNFRSF5 Protein (Cat.RP01218) was greater than 90% as determined by SEC-HPLC.



Immobilized Recombinant Human CD40 Ligand at 2µg/mL (100 µL/well) can bind Recombinant Human CD40 with a linear range of 224-898 ng/mL.