ABclonal www.abclonal.com

Recombinant Human IL-12/IL-12A&IL-12B Protein

Catalog No.: RP01232S Recombinant

Sequence Information

 Species
 Gene ID
 Swiss Prot

 HEK293 cells 3592&3593
 P29459&P2

 9460
 9460

Tags

C-His

Synonyms

CLMF;IL12;p70;Interleukin-12;IL12A;P3 5;NKSF

Product Information

Source Purification

Endotoxin

 $< 0.01EU/\mu g$ of the protein by LAL method.

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

Basic Information

Description

Recombinant Human IL-12/IL-12A&IL-12B Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (IL12B(Ile23-Ser328)-3 (GGGGS) IL12A(Arg23-Ser219)) of Human IL-12/IL-12A&IL-12B (Accession #) fused with a His tag at the C-terminus.

Bio-Activity

Measured by its ability to induce Interferon gamma secretion by human natural killer lymphoma NK-92 cells. The ED₅₀ for this effect is 0.94-3.75 ng/mL, corresponding to a specific activity of $2.67 \times 10^5 \sim 1.06 \times 10^6$ units/mg.

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at $2-8^{\circ}$ C for up to 1 week.

Avoid repeated freeze/thaw cycles.

^{*} For your safety and health, please wear a lab coat and disposable gloves when handling.

Validation Data



Recombinant Human IL-12/IL-12A&IL-12B Protein induce Interferon gamma secretion by human natural killer lymphoma NK-92 cells. The ED $_{50}$ for this effect is 0.94-3.75 ng/mL, corresponding to a specific activity of $2.67 \times 10^5 \sim 1.06 \times 10^6$ units/mg.