

Recombinant SARS-CoV-2 Spike S1(D614G) Protein

Catalog No.: RP01287 Recombinant 1 Publications

Sequence Information

Species Gene ID Swiss Prot HEK293 cells 43740568 P0DTC2

Tags

C-His

Synonyms

Envelope; SARS-CoV-2 Spike RBD (N501Y);Spike;Spike ECD;Spike RBD;Spike S1;Spike S2;Spike S2 ECD;S1-RBD protein;NCP-CoV RBD Protein; novel coronavirus RBD Protein;2019-nCoV RBD Protein;S glycoprotein Subunit1 RBD Protein

Product Information

Purification Source HEK293 cells > 95% by SDS-

PAGE.

Endotoxin

< 1.0 EU/µg of the protein by LAL method.

Formulation

Supplied as a 0.22 µm filtered solution in PBS, pH 7.4 or Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

Basic Information

Description

Recombinant SARS-CoV-2 Spike S1(D614G) Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Val16-Arg685(Asp614Gly)) of sars-cov-2 Spike S1(D614G) (Accession #QHD43416.1) fused with a 6×His tag at the C-terminus.

Bio-Activity


Measured by its binding ability in a functional ELISA. Immobilized recombinant SARS-CoV-2 Spike S1 at 2 µg/mL (100 µL/well) can bind recombinant Human ACE2 with a linear range of 0.03-6.26 ng/mL.

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.

Validation Data



Recombinant SARS-CoV-2 Spike S1(D614G) Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 110-130 kDa.



Immobilized recombinant SARS-CoV-2 Spike S1 at $2\mu g/mL$ (100 $\mu L/well$) can bind recombinant Human ACE2 with a linear range of 0.03-6.26 ng/mL.