ABclonal www.abclonal.com

Recombinant Human CCL17/TARC Protein

Catalog No.: RP01374 Recombinant

Sequence Information

Species Gene ID Swiss Prot HEK293 cells 6361 Q92583

Tags C-hFc&His

Synonyms

CCL17;A-152E5.3;ABCD-2;SCYA17;TAR C

Product Information

Source Purification
HEK293 cells > 90% by SDS-

PAGE.

Endotoxin

<0.1EU/µg

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

•

www.abclonal.com

Background

There are four members of the chemokine family: C-C kemokines, C kemokines, CXC kemokines and CX3C kemokines. The C-C kemokines have two cysteines nearby the amino terminus. There have been at least 27 distinct members of this subgroup reported for mammals, called C-C chemokine ligands-1 to 28. Chemokin ligand 17 (CCL17), also known as thymus and activation regulated chemokine(TARC), is a small cytokine belonging to the C-C chemokine family. CCL17 is expressed maily in thymus and transiently in phytohemagglutininstimulated peripheral blood mononuclear cells. CCL17 can induce chemotaxis in T cells by binding with the chemokine receptor CCR4.

Basic Information

Description

Recombinant Human CCL17/TARC Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ala24-Ser94) of human CCL17/TARC (Accession #NP_002978.1) fused with a Fc, 6×His tag at the C-terminus.

Bio-Activity

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

^{*} For your safety and health, please wear a lab coat and disposable gloves when handling.

Validation Data



Recombinant Human CCL17/TARC Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 38-40kDa.