

RP01382

Leader in Biomolecular Solutions for Life Science



Recombinant Human IL-1R3/IL-1RAcP/IL1RAP Protein

Catalog No.: RP01382 **Recombinant**

Sequence Information

Species **Gene ID** **Swiss Prot**
HEK293 cells 3556 Q9NPH3-1

Tags

C-hFc

Synonyms

IL1RAP;C3orf13;IL-1RAcP;IL1R3

Product Information

Source **Purification**
HEK293 cells > 90% by SDS-PAGE.

Endotoxin

<0.1EU/μg

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact



www.abclonal.com

Background

Interleukin-1 receptor accessory protein (IL-1RAcP) also known as Interleukin-1 receptor member 3 (IL-1R3) is a cytokine receptor that binds interleukin 1. The IL-1 receptor accessory protein (IL1RAP) is a transmembrane protein that interacts with IL-1R and is required for IL-1 signal transduction. Interleukin 1 induces the synthesis of the acute phase and proinflammatory proteins during infection, tissue damage, or stress, by forming a complex at the cell membrane with an interleukin 1 receptor and an accessory protein. IL-1RAcP/IL-1R3 is a necessary part of the interleukin 1 receptor complex which initiates signaling events that result in the activation of interleukin 1-responsive genes. Alternative splicing of this gene results in two transcript variants encoding two different isoforms, one membrane-bound and one soluble. The ratio of soluble to membrane-bound forms increases during acute-phase induction or stress. IL-1RAcP/IL-1R3 mediates interleukin-1-dependent activation of NF-kappa-B. Isoform 1 is part of the membrane-bound form of the IL-1 receptor. Signaling involves the formation of a ternary complex containing IL1R1, TOLLIP, MYD88, and IRAK1 or IRAK2. Isoform 2 modulates the response to interleukins by associating with soluble IL1R1 and enhancing interleukin-binding to the decoy receptor.

Basic Information

Description

Recombinant Human IL-1R3/IL-1RAcP/IL1RAP Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ser21-Glu359) of human IL1RAP/IL1R3 (Accession #NP_002173.1) fused with a hFc tag at the C-terminus.

Bio-Activity

Storage


Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.


Avoid repeated freeze/thaw cycles.

* For your safety and health, please wear a lab coat and disposable gloves when handling.

Validation Data



Recombinant Human IL-1R3/IL-1RAcP/IL1RAP Protein was determined by SDS-PAGE with Coomassie Blue, showing a band at 90-100kDa.



The purity of Human IL1RAP/IL1R3 Protein (Cat.RP01382) was greater than 90% as determined by SEC-HPLC.