

RP01665

Leader in Biomolecular Solutions for Life Science



Recombinant Mouse Thrombopoietin/THPO Protein

Catalog No.: RP01665

Recombinant

Sequence Information

Species	Gene ID	Swiss Prot
HEK293 cells	21832	P40226-1

Tags

C-6His

Synonyms

MI; Tpo; Mgdf; Mpllg; THPO; TPO

Product Information

Source	Purification
HEK293 cells	> 95% by SDS-PAGE.

Endotoxin

<0.1EU/μg

Formulation

Lyophilized from a 0.22 μm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stabilizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Background

Thrombopoietin (TPO or THPO), also known as myeloproliferative leukemia virus ligand (c-Mpl), is a hematopoietic growth factor belonging to the EPO/TPO family. The thrombopoietin protein is produced mainly by the liver and the kidney that regulates the production of platelets by the bone marrow. Thrombopoietin protein stimulates both proliferation of progenitor megakaryocytes and their maturation to platelet-producing megakaryocytes, and also accelerates the recovery of platelets. Thrombopoietin protein is involved in cardiovascular disease as it regulates megakaryocyte development and enhances platelet adhesion/aggregation. It has been identified that surface c-MPL, the receptor for thrombopoietin protein, binds to the ligand and mediates the action.

Basic Information

Description

Recombinant Mouse Thrombopoietin/THPO Protein is produced by HEK293 cells expression system. The target protein is expressed with sequence (Ser22-Thr356) of mouse Thrombopoietin/THPO (Accession #NP_033405.1.) fused with and a 6×His tag at the C-terminus.

Bio-Activity

Measured in a cell proliferation assay using MO7e human megakaryocytic leukemic cells. Avanzi, G. et al. (1988) Br. J. Haematol. 69:359. The ED₅₀ for this effect is 0.45-1.80 ng/mL.

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt. After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week. Avoid repeated freeze/thaw cycles.


Contact




www.abclonal.com

* For your safety and health, please wear a lab coat and disposable gloves when handling.

Validation Data



Recombinant Mouse Thrombopoietin/THPO Protein was determined by SDS-PAGE under reducing conditions with Coomassie Blue.



Recombinant Mouse THPO stimulates cell proliferation of the MO7e human megakaryocytic leukemic cells.The ED₅₀ for this effect is 0.45-1.80 ng/mL.