ABclonal www.abclonal.com

Recombinant Human FGF-4 Protein

Catalog No.: RP01712 Recombinant

Sequence Information

 Species
 Gene ID
 Swiss Prot

 <I>E.
 2249
 P08620-1

 coli </I>
 Policy II
 Policy II

Tags NO-tag

Synonyms

HST; KFGF; FGF-4; HST-1; HSTF1; K-FGF; HBGF-4; HSTF-1;FGF4

Product Information

Source Purification <I>E. coli</I> > 95% by SDS-

PAGE.

Endotoxin

< 0.1EU/µg

Formulation

Lyophilized from a 0.22 µm filtered solution of PBS, pH 7.4.

Reconstitution

Centrifuge the vial before opening. Reconstitute to a concentration of 0.1-0.5 mg/mL in sterile distilled water. Avoid vortex or vigorously pipetting the protein. For long term storage, it is recommended to add a carrier protein or stablizer (e.g. 0.1% BSA, 5% HSA, 10% FBS or 5% Trehalose), and aliquot the reconstituted protein solution to minimize free-thaw cycles.

Contact

•

www.abclonal.com

Background

FGF (fibroblast growth factor) signalling is known to be required for many aspects of mesoderm formation and patterning during Xenopus development and has been implicated in regulating genes required for the specification of both blood and skeletal muscle lineages. Fibroblast growth factor 4 (FGF4) signaling induces differentiation from embryonic stem cells (ESCs) via the phosphorylation of downstream molecules such as mitogen-activated protein kinase/extracellular signal-related kinase (MEK) and extracellular signal-related kinase 1/2 (ERK1/2). Fibroblast Growth Factor 4 (FGF-4) could not only increase the proliferation of bone marrow mesenchymal stem cells (BMSCs), but also induce BMSCs into hepatocyte-like cells in vitro. FGF4 transduced BMSCs contributed to liver regeneration might by the transplanted microenvironment. The FGF4-bFGF BMSCs thus can enhance the survival of the transplanted cells, diminish myocardial fibrosis, promote myocardial angiogenesis, and improve cardiac functions.

Basic Information

Description

Recombinant Human FGF-4 Protein is produced by *E. coli* expression system. The target protein is expressed with sequence (Gly25-Leu206) of human FGF-4 (Accession #NP_001998.1) fused with no additional amino acid.

Bio-Activity

Storage

Store at -20°C. Store the lyophilized protein at -20°C to -80 °C up to 1 year from the date of receipt.

After reconstitution, the protein solution is stable at -20°C for 3 months, at 2-8°C for up to 1 week.

Avoid repeated freeze/thaw cycles.

^{*} For your safety and health, please wear a lab coat and disposable gloves when handling.