

ZNF225 Ab

Cat.#: AF0511
Size: 100ul,200ul

Concn.: 1mg/ml
Source: Rabbit

Mol.Wt.: 82kDa
Clonality: Polyclonal

Application: WB 1:500-1:2000, IF/ICC 1:100-1:500

Reactivity: Human

Purification: The antiserum was purified by peptide affinity chromatography using SulfoLink™ Coupling Resin (Thermo Fisher Scientific).

Specificity: ZNF225 Ab detects endogenous levels of ZNF225.

Immunogen: A synthesized peptide derived from human ZNF225.


Uniprot: Q9UK10

Description: ZNF225 May be involved in transcriptional regulation.


Subcellular Location: Nucleus.

Similarity: Belongs to the krueppel C2H2-type zinc-finger protein family.

Storage Condition and Buffer: Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of extracts from Colo205, using ZNF225 Ab. The lane on the left was treated with blocking peptide.



Western blot analysis on HeLa cell lysate using ZNF225 Ab, The lane on the left is treated with the antigen-specific peptide.



AF0511 staining HeLa by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.