

MAPKAPK5 Ab

Cat.#: AF0835 Concn.: 1mg/ml Mol.Wt.: 60kDa Size: 100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IF/ICC 1:100-1:500

Reactivity: Human, Mouse

Purification: The antiserum was purified by peptide affinity

chromatography using SulfoLink™ Coupling Resin (Thermo

Fisher Scientific).

Specificity: MAPKAPK5 Ab detects endogenous levels of MAPKAPK5.

Immunogen: A synthesized peptide derived from human MAPKAPK5.

Uniprot: Q8IW41

Description: The protein encoded by this gene is a member of the

serine/threonine kinase family. In response to cellular stress and proinflammatory cytokines, this kinase is activated through its phosphorylation by MAP kinases including MAPK1/ERK, MAPK14/p38-alpha, and MAPK11/p38-beta. In vitro, this kinase phosphorylates heat shock protein HSP27 at its physiologically relevant sites. Two alternately spliced transcript variants of this gene encoding distinct isoforms

have been reported.

Subcellular Location: Cytoplasm. Nucleus. Also observed in the nucleus.

Tissue Specificity: Expressed ubiquitously.

Similarity: Belongs to the protein kinase superfamily. CAMK Ser/Thr


protein kinase family.

Storage Condition and

Buffer:

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20

°C.Stable for 12 months from date of receipt.



Western blot analysis on K562 cell lysate using MAPKAPK5 Ab,The lane on the left is treated with the antigen-specific

peptide.



Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com



AF0835 staining Hela by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween020 at 4° C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.