Phospho-CENP-A (Ser7) Ab

Cat.#: AF2330 Concn.: 1mg/ml Mol.Wt.: 17kDa Size: 100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IHC 1:50-1:200

Reactivity: Human,Rat

Purification: The Ab is from purified rabbit serum by affinity purification

via sequential chromatography on phospho- and non-

phospho-peptide affinity columns.

Specificity: Phospho-CENP-A (Ser7) Ab detects endogenous levels of

CENP-A.

Immunogen: A synthesized peptide derived from human CENP-A around

the phosphorylation site of Ser7.

Uniprot: P49450

Subcellular Location: Nucleus. Chromosome > centromere > kinetochore.

Localizes exclusively in the kinetochore domain of centromeres. Occupies a compact domain at the inner kinetochore plate stretching across 2 thirds of the length of the constriction but encompassing only one third of the

constriction width and height.

Tissue Specificity: Expression varies across the cell cycle, with high levels in G2

phase (at the mRNA level).

Similarity: The CATD (CENPA targeting domain) region is responsible

for the more compact structure of nucleosomes containing CENPA (PubMed:15282608). It is necessary and sufficient to

mediate the localization into centromeres

(PubMed:7962047, PubMed:15282608).Belongs to the

histone H3 family.

Storage Condition and

Buffer:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20

°C.Stable for 12 months from date of receipt.



Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com



Western blot analysis of extracts of HeLa , using Phospho-CENP-A (Ser7) Ab.



AF2330 at 1/100 staining Rat liver tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1% TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.