

## Phospho-MFF (Ser172) Ab

Cat.#: AF2365  
 Size: 100ul,200ul

Concn.: 1mg/ml  
 Source: Rabbit

Mol.Wt.: 25, 27kDa  
 Clonality: Polyclonal

**Application:** WB 1:500-1:2000, IHC 1:50-1:200

**Reactivity:** Human

**Purification:** The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.

**Specificity:** Phospho-MFF (Ser172) Ab detects endogenous levels of MFF.

**Immunogen:** A synthesized peptide derived from human MFF around the phosphorylation site of Ser172.


**Uniprot:** Q9GZY8

**Subcellular Location:** Mitochondrion outer membrane.


**Tissue Specificity:** Highly expressed in heart, kidney, liver, brain, muscle, and stomach.

**Similarity:** Belongs to the Tango11 family.

**Storage Condition and Buffer:** Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of extracts of MCF-7 cells, using Phospho-MFF (Ser172) Ab.



AF2365 at 1/100 staining Human thyroid cancer tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.

**IMPORTANT:** For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.