

Phospho-Chk1 (Ser345) Ab

Cat.#: AF3008 Mol.Wt.: kDa Concn.: 1mg/ml

Size: 100ul.200ul. Source: Rabbit Clonality: Polyclonal

Application: IHC 1:50-1:200. IF/ICC 1:100-1:500

Reactivity: Human, Mouse, Rat

Purification: The Ab is from purified rabbit serum by affinity purification

via sequential chromatography on phospho- and non-

phospho-peptide affinity columns.

Specificity: Phospho-Chk1 (Ser345) Ab detects endogenous levels of

Chk1 only when phosphorylated at Serine 345.

A synthesized peptide derived from human Chk1 around the Immunogen:

phosphorylation site of Serine 345.

Uniprot: 014757

Description: DNA damage induced protein phosphorylation; regulation of

mitotic centrosome separation; regulation of S phase; peptidyl-threonine phosphorylation: DNA repair: chromatinmediated maintenance of transcription; negative regulation

of mitosis:

Cvtoskeleton; Nucleus; Subcellular Location:

Tissue Specificity: Expressed ubiquitously with the most abundant expression

in thymus, testis, small intestine and colon.

Similarity: The autoinhibitory region (AIR) inhibits the activity of the

> kinase domain. Belongs to the protein kinase superfamily. CAMK Ser/Thr protein kinase family. NIM1 subfamily.

Storage Condition and

Buffer:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20

°C.Stable for 12 months from date of receipt.



Phospho-Chk1 (Ser345) Ab for IHC in human brain.



Affinity Biosciences website:www.affbiotech.com



AF3008 staining HeLa by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween®20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.