


Phospho-Abl (Tyr204) Ab

Cat.#: AF3038
Size: 100ul,200ul


Concn.: 1mg/ml
Source: Rabbit

Mol.Wt.: 135kDa
Clonality: Polyclonal


Application:	WB 1:500-1:2000, IF/ICC 1:100-1:500
Reactivity:	Human,Mouse,Rat,Monkey
Purification:	The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Specificity:	Phospho-Abl (Tyr204) Ab detects endogenous levels of Abl only when phosphorylated at Tyrosine 204.
Immunogen:	A synthesized peptide derived from human Abl around the phosphorylation site of Tyrosine 204.
Uniprot:	P00519
Description:	The ABL1 protooncogene encodes a cytoplasmic and nuclear protein tyrosine kinase that has been implicated in processes of cell differentiation, cell division, cell adhesion, and stress response. Activity of c-Abl protein is negatively regulated by its SH3 domain, and deletion of the SH3 domain turns ABL1 into an oncogene.
Subcellular Location:	Cytoplasm > cytoskeleton. Nucleus. Sequestered into the cytoplasm through interaction with 14-3-3 proteins and Nucleus membrane. The myristoylated c-ABL protein is reported to be nuclear.
Tissue Specificity:	Widely expressed.
Similarity:	Belongs to the protein kinase superfamily. Tyr protein kinase family. ABL subfamily.
Storage Condition and Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of extracts from HepG2, using Phospho-Abl (Tyr204) Ab. Lane 1 was treated with the blocking peptide.



Western blot analysis of Phospho-Abl (Tyr204) Ab expression in Adriamycin treated COS7 cells lysates. The lane on the right is treated with the antigen-specific peptide.



Western blot analysis of Abl phosphorylation expression in Adriamycin treated COS7 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



AF3038 staining COS7 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.