

Phospho-PKCB (Ser661) Ab

Cat.#: AF3096 Concn.: 1mg/ml Mol.Wt.: 77kDa Size: 100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000, IF/ICC 1:100-1:500

Reactivity: Human, Mouse, Rat

Purification: The Ab is from purified rabbit serum by affinity purification

via sequential chromatography on phospho- and non-

phospho-peptide affinity columns.

Specificity: Phospho-PKCB (Ser661) Ab detects endogenous levels of

PKCB only when phosphorylated at Serine 661.

Immunogen: A synthesized peptide derived from human PKCB around the

phosphorylation site of Serine 661.

Uniprot: P05771

Description: Pim1 is thought to play a role in signal transduction in blood

cells. May affect the structure or silencing of chromatin by

phosphorylating HP1 gamma/CBX3.

Subcellular Location: Cytoplasm. Nucleus. Membrane.

Similarity: Belongs to the protein kinase superfamily. AGC Ser/Thr


protein kinase family. PKC subfamily.

Storage Condition and

Buffer:

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20

°C.Stable for 12 months from date of receipt.



Western blot analysis of Phospho-PKCB (Ser661) Ab expression in heatshock treated Hela cells lysates. The lane on the right is treated with the antigen-specific peptide.



Affinity Biosciences website:www.affbiotech.com

website:www.affbiotech.com order:order@affbiotech.com



Western blot analysis of PKCB phosphorylation expression in heatshock treated HeLa whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



AF3096 staining HeLa by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100,then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1% TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.