

Interleukin 12A Ab

Cat.#: AF5133 Concn.: 1mg/ml Mol.Wt.: 24 kDa Size: 100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000 IHC 1:50-1:200

Reactivity: Human, Mouse, Rat

Purification: The antiserum was purified by peptide affinity

chromatography using SulfoLink™ Coupling Resin (Thermo

Fisher Scientific).

Specificity: Interleukin 12A Ab detects endogenous levels of total

Interleukin 12A.

Immunogen: A synthesized peptide derived from human Interleukin 12A.

Uniprot: P29459

Description: Cytokine that can act as a growth factor for activated T and

NK cells, enhance the lytic activity of NK/lymphokineactivated Killer cells, and stimulate the production of IFN-

gamma by resting PBMC.

Subcellular Location: Secreted

Tissue Specificity: Down-regulated in response to enterovirus 71 (EV71)

infection.


Similarity: Belongs to the IL-6 superfamily.

Storage Condition and

Buffer:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20

°C.Stable for 12 months from date of receipt.



Western blot analysis of Interleukin 12A expression in HeLa cells, The lane on the left is treated with the antigen-specific


peptide.



Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com



AF5133 at 1/100 staining Human Melanoma tissue by IHC-P. The sample was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The sample was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.



AF5133 at 1/100 staining human kidney tissues sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at127

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1% TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.