PERFORMANCE DATA SHEET

1751

Monoclonal anti-human CD11a*

mAb name/Clone: 38 Isotype: Mouse IgG2a

Immunogen: Human monocytes

CATALOG#: 158-820 (Preservative Free)

QUANTITY: 100 μg CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD11a (α L integrin) complexes with CD18 (β_2 integrin) to form the LFA-1 heterodimer adhesion molecule which binds to three ICAM's 1, 2, and 3, also designated CD54, CD102, and CD50 respectively. LFA-1 is expressed on lymphocytes, monocytes and neutrophils and plays a role in cell-cell adhesion. Antibody 38 recognizes the I domain of the integrin alpha L subunit (CD11a) of about 180 kd. Antibody 38 blocks binding of ICAM-1 and ICAM-3 to LFA-1 at 5-10 µg/ml.

References: I. Dransfield & N. Hogg, (1989) EMBO J **12**: 3759-3765. Leukocyte Typing V (S.F. Schlossman,et al, eds.) Oxford University Press, Oxford, (1995) p. 1581-1592. R.C. Landis, et al, (1994) J Cell Biol **126**: 529-537. C.L. Holness, et al, (1995) J Biol Chem **270**: 877-884.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended. Open under aseptic conditions.


PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:_____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl.

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE with less than 1% Bovine Immunoglobulin. Product was 0.2μ sterile filtered and vialed under aseptic conditions.

PERFORMANCE: Five x 10^5 cultured **HPB-MLT** human tumor cells were washed and incubated 45 minutes on ice with 80 µl of anti-CD11a antibody at **5** µg/ml. Cells were washed twice and incubated with 2^0 reagent Goat anti-Mouse IgG/FITC (Catalog #232-011), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **1.28** log₁₀ fluorescent units when compared to a Mouse IgG2a negative control (Catalog #281-010) at a similar concentration.

Binding of anti-CD11a mAb + GAM/FITC to human HPB-MLT cells



*This Product is intended for Laboratory Research use only.