PERFORMANCE DATA SHEET

1818

Monoclonal anti-human CD100*

mAb name/Clone: 133-1C6 *Isotype:* Mouse IgM

Immunogen: PHA stimulated human PBL

CATALOG#: 321-820 (Preservative-free) LOT#: 634003

QUANTITY: 100 μg CONCENTRATION: 1.0 mg/ml

INFORMATION: Human CD100 is a transmembrane glycoprotein that belongs to the semaphorin gene family and is found on most hematopoietic cells except RBC, platelets and immature bone marrow cells. CD100 plays a key role in lymphocyte activation (2, 3). Antibody 133-1C6 recognizes a CD100 molecule of about 150 kd. *References:* 1)Leukocyte Typing VI (T. Kishimoto, et al, eds.) Garland Publishing, Inc., New York (1997) p. 534-539. 2)K.T. Hall, et al, (1996) Proc Natl Acad Sci USA 93:11780-11785. 3)A. Elhabazi, et al, (1997) J Biol Chem 272: 23515-23520.

STORAGE CONDITIONS: Store at 2 - 5°C. Freeze/Thawing is not recommended. Open under aseptic conditions.

PRODUCT STABILITY: Product should retain activity for at least 12 months after shipping date when stored as recommended. Ship Date:_____

BUFFER: 50 mM Sodium Phosphate pH 7.5, 100 mM Potassium Chloride, 150mM NaCl.

PRODUCTION: Antibody was Protein A purified from (low FBS containing) tissue culture supernatant. Purity was >95% Immunoglobulin by SDS-PAGE and contains less than 1% Bovine Immunoglobulin. Product was 0.2μ sterile filtered and vialed under aseptic conditions.

PERFORMANCE: Five x 10^5 cultured human **Jurkat** cells were incubated 45 minutes on ice with 80 μ l of anti-CD100 antibody at **10** μ g/ml. Cells were washed twice and incubated with 2^o reagent Goat anti-Mouse IgG/FITC (Catalog #232-011; this reagent cross reacts well with Mouse IgM), after which they were washed three times, fixed and analyzed by FACS. Cells stained positive with a mean shift of **0.82** \log_{10} fluorescent units when compared to a Mouse IgM negative control (Catalog #290-010) at a similar concentration.

*This Product is intended for Laboratory Research use only.

Binding of anti-CD100 antibody to human cell lines

