

Rabbit Anti-Human VEGF-A

ORDERING INFORMATION

Catalog Number:	102-PA03AG
Size:	50 µg
Formulation:	Polyclonal Antibody ; Lyophilized
Synonyms:	VEGFA; VPF; VEGF; MVCD1
Antigen:	Recombinant human VEGF165 RT #300-036)
Application:	WB, IP
Stabilizer	None
Buffer:	PBS pH 7.4 w/o preservative

Description:

Vascular endothelial growth factor (VEGF or VEGF-A), also known as vascular permeability factor (VPF) or vasculotropin, is a homodimeric 34 - 42 kDa, heparin-binding glycoprotein with potent angiogenic, mitogenic and vascular permeability-enhancing activities specific for endothelial cells. Different isoforms can be generated by differential splicing (e.g. VEGF165). All eight cysteine residues involved in intra- and inter-chain disulfide bonds are conserved among these growth factors. A cDNA encoding a protein having a 53% amino acid sequence homology in the PDGF-like region of VEGF has been isolated from a human placental cDNA library. This protein, named placenta growth factor (PlGF), is now recognized to be a member of the VEGF family of growth factors. Two receptor tyrosine kinases have been described as putative VEGF receptors. Flt-1 (fms-like tyrosine kinase), and KDR (kinase-insert-domain-containing receptor) proteins have been shown to bind VEGF-A with high affinity. In vitro, VEGF is a potent endothelial cell mitogen. The antibody is against the N-terminus and will recognize all VEGF-A isoforms.

Reconstitution:

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Stability:

The lyophilized antibody is stable at room temperature for up to 1 month. The reconstituted antibody is stable for at least two weeks at 2-8 °C. Frozen aliquots are stable for at least 6 months when stored at -20 °C. **Avoid repeated freeze-thaw cycles!**

Optimal dilutions should be determined by each laboratory for each application.

The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users!

This product is sold for Research Use Only !

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Tel: 617-549-2665; Fax: (480) 247-4337, angioproteomie@gmail.com