

## Rabbit Anti-Rat VEGF-C

#### **ORDERING INFORMATION**

Catalog Number: 102-PA03S

Size:  $100 \,\mu g$ 

Formulation: Polyclonal Antibody; Lyophilized

**Synonyms:** VEGF C

**Antigen:** Recombinant rat VEGF-C (RT #R20-015)

**Application:** WB, E, IP

**Stabilizer** None

**Buffer:** PBS pH 7.4 w/o preservative

## Description:

Vascular endothelial growth factor (VEGF or VEGF-A), also known as vascular permeability factor (VPF) or vasculotropin, is a homodimeric 34 - 42 kDa, heparin-binding glycoprotein with potent angiogenic, mitogenic and vascular permeability-enhancing activities specific for endothelial cells. Different isoforms can be generated by differential splicing (e.g. VEGF165). All eight cysteine residues involved in intra- and inter-chain disulfide bonds are conserved among these growth factors. A cDNA encoding a protein having a 53% amino acid sequence homology in the PDGF-like region of VEGF has been isolated from a human placental cDNA library. This protein, named placenta growth factor (PlGF), is now recognized to be a member of the VEGF family of growth factors. Two receptor tyrosine kinases have been described as putative VEGF receptors. Flt-1 (fms-like tyrosine kinase), and KDR (kinase-insert-domain-containing receptor) proteins have been shown to bind VEGF-A with high affinity. In vitro, VEGF is a potent endothelial cell mitogen. The antibody is against the N-terminus and will recognize all VEGF-A isoforms.

#### Reconstitution:

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

## Stability:

The lyophilized antibody is stable at room temperature for up to 1 month. The reconstituted antibody is stable for at least two weeks at 2-8 °C. Frozen aliquots are stable for at least 6 months when stored at -20 °C. **Avoid repeated freeze-thaw cycles!** 

Optimal dilutions should be determined by each laboratory for each application.

The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users!

# This product is sold for Research Use Only!

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Tel: 617-549-2665; Fax: (480) 247-4337, angioproteomie@gmail.com