

## Rabbit Anti-Human VEGFR-3/FLT-4

#### **ORDERING INFORMATION**

Catalog Number: 102-PA22S

Size:  $100 \mu g$ 

Formulation: Polyclonal Antibody; Lyophilized

Synonyms: FLT4; PCL; LMPH1A; VEGFR3

Antigen: RH sFLT-4 (D1-7) (RT #S01-017)

**Application:** WB, IP, E, IHC (C)

Label: None

**Buffer:** PBS pH 7.4 w/o preservative

### Description:

Receptor tyrosine Kinase VEGFR-3, also known as FLT4, together with VEGFR1 (FIT1) and VEGFR2 (KDR/Flk-1), are the receptors for vascular endothelial growth factors (VEGF). The VEGFR family belongs to the class II subfamily of receptor tyrosine kinases (RTKs), containing a large extracellular region which is composed of seven Ig-like domains (D1–D7), a single transmembrane (TM) helix and cytoplasmic region with a tyrosine kinase activity. In VEGFR-3, the fifth Ig homology domain is proteolytically cleaved which results in polypeptides remain linked by two disulfide bonds. VEGFR-3 is widely expressed on all endothelia cells in early embryogenesis, while, in adult tissues, VEGFR-3 expression disappears from the vascular endothelial cells and is observed only on the lymphatic endothelium. VEGF-C and VEGF-D activation of VEGFR-3 plays an important role in the formation of the lymphatic vessel system.

#### Reconstitution:

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

### Stability:

The lyophilized antibody is stable at room temperature for up to 1 month. The reconstituted antibody is stable for at least two weeks at 2-8 °C. Frozen aliquots are stable for at least 6 months when stored at -20 °C. **Avoid repeated freeze-thaw cycles!** 

Optimal dilutions should be determined by each laboratory for each application.

The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users!

# This product is sold for Research Use Only!