

Rabbit Anti-Rat VEGF-C

ORDERING INFORMATION

Catalog Number:	104-PA10S
Size:	100 µg
Formulation:	Polyclonal Antibody ; Lyophilized
Synonyms:	vascular endothelial growth factor C
Antigen:	N-terminal peptide
Application:	WB, E, IP
Stabilizer	None
Buffer:	PBS pH 7.4 w/o preservative

Description:

VEGF-C, a member of the VEGF/PDGF family of structurally related proteins, is a potent angiogenic cytokine. It promotes endothelial cell growth, promotes lymphangiogenesis, and can also affect vascular permeability. VEGF-C is expressed in various tissues, but is not produced in peripheral blood lymphocytes. It forms cell surfaced-associated non-covalent disulfide linked homodimers, and can bind and activate both VEGFR-2 (flk1) and VEGFR-3 (flt4) receptors. During embryogenesis, VEGF-C may play a role in the formation of the venous and lymphatic vascular systems. Both VEGF-C and VEGF-D are over-expressed in certain cancers, and the resulting elevated levels of VEGF-C or VEGF-D tend to correlate with increased lymphatic metastasis. Recombinant, fully processed rat VEGF-C is a 16-18 kDa non-disulfide linked homodimeric protein consisting of two 116 amino acid polypeptide chains. Due to glycosylation the protein migrates as a 20.0-22.0 kDa band under non-reducing condition.

Reconstitution:

Centrifuge vial prior to opening. Reconstitute in sterile water to a concentration of 0.1-1.0 mg/ml.

Stability:

The lyophilized antibody is stable at room temperature for up to 1 month. The reconstituted antibody is stable for at least two weeks at 2-8 °C. Frozen aliquots are stable for at least 6 months when stored at -20 °C. **Avoid repeated freeze-thaw cycles!**

Optimal dilutions should be determined by each laboratory for each application.

The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users!

This product is sold for Research Use Only !