


Rabbit anti p38 MAPK (Ab-182)

- Catalog Number:** AP21245
Amount: 100µl (for 10 Western Blotting)
Swiss-Prot No. : Q16539
Form of Antibody: Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage/Stability: Store at -20°C/1 year
Immunogen: The antiserum was produced against synthesized non-phosphopeptide derived from human P38MAPK around the phosphorylation site of tyrosine 182 (T-G-Y^P-V-A).
Purification: The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Specificity/Sensitivity P38MAPK (Ab-182) antibody detects endogenous levels of total P38MAPK protein.
Reactivity: Human, Mouse, Rat
Applications: WB: 1:500~1:1000
References: Ming Zheng, et al.(2005) The FASEB Journal. 19: 109-111
 Bernt van den et al.(2001) Blink Immunology, 166: 582-587
 Arshad Rahman, et al. (2004) Am J Physiol Lung Cell Mol Physiol 287: L1017-L1024
 Osamu Yoshino, et al. (2003) Endocrinology & Metabolism Vol. 88: 2236-2241



Western blot analysis of extracts from NIH-3T3 (Line 1, 4, 7 and 10) and cos7 (Line 2, 5, 8 and 11 and K562 (Line 3, 6, 9 and 12) cells, untreated or treated with UV (20min), using P38 MAPK (Ab-182) antibody (AP21245, Lane 1, 2, 3, 4, 5 and 6) and P38 MAPK (phospho- Tyr182) antibody (pAP11253, Lane 7, 8, 9, 10, 11 and 12).

For Research Use Only

Contact & Ordering Information: Angio-Proteomie, 11 Park Drive, Suite 12, Boston, MA 02215, USA. Fax: (480) 247-4337, angioproteomie@gmail.com