

GFP Expressing Human Ovarian Cancer Cells (SK-OV-3)
ORDER INFORMATION

Name of Cells: GFP Expressing Human Ovarian Cancer Cells (GFP-SK-OV-03)
Catalogue Number: **cAP-0054GFP**
Product Format: Frozen Vial
Cell Number: >5 x 10⁵/vial

General Information

Human SK-OV-3 Cells was established from the ascites of a 64-year-old Caucasian female with adenocarcinoma of the ovary. GFP-Expressing SK-OV-3 cells are selected from SK-OV-3 (cAP-0054) transfected with GFP expressing lentiviruses resistant to puromycin. The cells are supplied in frozen vials with more than 5 x 10⁵ cells/vial. Universal Full Growth Medium (cAP-01B) is recommended to culture the cells.


Human SK-OV-3 cells are negative for HIV-1, HBV, HCV, and mycoplasma.

Product Use: GFP Expressing Human SK-OV-3 cells are for Research Use Only.

Shipping: Frozen Vial.

Handling of Arriving Cells

When you receive the cells in a frozen vial, you can transfer the vial of cells into a -80°C freezer for short period storage or a liquid nitrogen tank for long term storage. Thaw the cells in a 37°C water bath, and then transfer the cells in a T25 flask in 5 ml of Universal Full Growth Medium (cAP-01B).



GFP Expressing SK-OV-3 Cells

Subculture Protocol

- A) Rinse the cells in T25 flask with 5ml HBSS (**Room Temperature, RT**) twice.
- B) Add 2ml of Trypsin/EDTA (**RT**) (cAP-23) into one T25 flask (make sure the whole surface of the T25 flask is covered with Trypsin/EDTA), and gently dispose the excessive Trypsin/EDTA solution **within 20 seconds** with aspiration.
- C) Leave the T25 flask with the cells at **RT** for 1 minute (the cells usually will detach from the surface within 1-2 minutes). You can monitor the cells under microscope and when most of cells become rounded up, hit the flask against the bench surface, and the cells will move on the surface of the flask when monitoring under microscope.
- D) Add 5ml Trypsin Neutralization Buffer and spin the cells down with 800g for 5 minutes.
- E) Re-suspend the cell pellet with 15ml 10%FBS in DMEM and the cell suspension is transferred directly into 3 T25 flasks (5ml each, and the cells are sub-cultured at 1:3 ratios)
- F) Change medium every 2-3 days and cells usually become confluent within 7-8 days.

Related products

Quick Coating Solution	cAP-01	240ml	Angio-Proteomie
Universal Full Growth Medium	cAP-01B	500ml	Angio-Proteomie
HBSS w/o Ca ²⁺ , Mg ²⁺	cAP-11	100ml	Angio-Proteomie
Cell Freezing Solution (FBS)	cAP-22	50ml	Angio-Proteomie
Cell Freezing Solution (Non-FBS)	cAP-22B	50ml	Angio-Proteomie
Trypsin/EDTA Solution	cAP-23	100ml	Angio-Proteomie
Trypsin Neutralization Solution	cAP-28	100ml	Angio-Proteomie
ITS (100x)	cAP-26	10ml	Angio-Proteomie
L-Glutamine-MAXIMUM (100x)	cAP-27	100ml	Angio-Proteomie
Human Plasma Fibronectin Solution	cAP-42	1mg/ml	Angio-Proteomie

Caution: Handling human tissue derived products is potentially bio-hazardous. Although each cell strain is tested negative for HIV, HBV and HCV DNA, diagnostic tests are not necessarily 100% accurate; therefore, proper precautions must be taken to avoid inadvertent exposure. Always wear gloves and safety glasses when working these materials. Never mouth pipette. We recommend following the universal procedures for handling products of human origin as the minimum precaution against contamination.