

**EA.hy926 Cells**
**ORDER INFORMATION**

**Name of Cells:** EA.hy926 Cells  
**Catalogue Number:** **cAP-0059**  
**Product Format:** Frozen Vials  
**Cell Number:** >5x 10<sup>5</sup>/vial

**General Information**

The human umbilical vein cell line, EA.hy926, was established by fusing primary human umbilical vein cells with a thioguanine-resistant clone of A549 by exposure to polyethylene glycol (PEG). Hybrid clones were selected in HAT medium and screened for factor VIII-related antigen.

Endothelial Growth Medium (EGM, contains 10% serum and growth supplements, cAP-02) is recommended for cell culture and these cells have a minimum average population doubling levels > **50** when cultured following the detailed protocol described below.

**Characterization of the cells**

The cells express von Willebrand's factor (vWF).

**Product Use:** EA.hy926 cells are for research use only.

**Shipping:** Frozen Vials.

**Handling of Arriving Cells**

When you receive the cells in a frozen vial, you can transfer the vial of cells into a -80°C freezer for short period storage or a liquid nitrogen tank for long term storage. Thaw the cells in a 37°C water bath, and then transfer the cells into a T25 flask pre-coated with Quick coating solution (cAP-01) as described in details in Subculture Protocol.

**Subculture Protocol**

- A) Pre-coating of T25 flasks: Add 2ml of Quick Coating Solution (**cAP-01**) into one T25 flask and make sure whole surface of the flask is covered with the coating solution. Five minutes later, dispose excessive Quick Coating Solution by aspiration and the flask is ready to be used (no need for overnight incubation when using Quick Coating Solution). Other extracellular matrix can be used including gelatin, collagen, and fibronectin and you are advised to test the conditions for using those materials in advance.
- B) Rinse the cells in T25 flask with 5ml HBSS (**Room Temperature, RT**) twice.
- C) Add 2ml of Trypsin/EDTA (**RT**) (cAP-23) into one T25 flask (make sure the whole surface of the T25 flask is covered with Trypsin/EDTA), and gently dispose the excessive Trypsin/EDTA solution **within 60 seconds** with aspiration.
- D) Leave the T25 flask with the cells at 37C for extra 1-2 minute (the cells usually will detach from the surface within 1-2 minutes). You can monitor the cells under microscope and when most of cells become rounded up, hit the flask against the bench surface, and the cells will move on the surface of the flask when monitoring under microscope.
- E) Add 5ml Trypsin Neutralization Buffer and spin the cells down with 800g for 5 minutes.
- F) Re-suspend the cell pellet with 10ml of HDaNC growth medium and the cell suspension is transferred directly into 2 pre-coated T25 flasks (5ml each, and the cells are sub-cultured at 1:2)
- G) Change medium every 2-3 days and cells usually become confluent within 7 days (when split at a 1:4 ratio).

**Related products**

Quick Coating Solution	cAP-01	240ml	Angio-Proteomie
Endothelial Growth Medium	cAP-02	500ml	Angio-Proteomie
HBSS w/o Ca <sup>2+</sup> , Mg <sup>2+</sup>	cAP-11	100ml	Angio-Proteomie
Trypsin/EDTA Solution	cAP-23	100ml	Angio-Proteomie
Trypsin Neutralization Solution	cAP-28	100ml	Angio-Proteomie

**Caution:** Handling human tissue derived products is potentially bio-hazardous. Although each cell strain is tested negative for HIV, HBV and HCV DNA, diagnostic tests are not necessarily 100% accurate; therefore, proper precautions must be taken to avoid inadvertent exposure. Always wear gloves and safety glasses when working these materials. Never mouth pipette. We recommend following the universal procedures for handling products of human origin as the minimum precaution against contamination.