

## Goat anti-FRAT2 / GSK-3 Antibody

<b>Item Number</b>	dAP-0083
<b>Target Molecule</b>	Principle Name: FRAT2 / GSK-3; Official Symbol: FRAT2; All Names and Symbols: FRAT2; GSK-3; frequently rearranged in advanced T-cell lymphomas 2; GSK-3 binding protein FRAT2; RP11-452K12.2; MGC10562; Accession Number (s): NP_036215.1; Human Gene ID(s): 23401; Non-Human GeneID(s):
<b>Immunogen</b>	GPDRIALQPSGSL, is from C Terminus
<b>Applications</b>	Pep ELISA, WB Species Tested: Human
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 2000.
<b>Western Blot</b>	Western Blot: Approx 30-35kDa band seen in Human Heart lysate. Recommended for use at 1-3µg/ml.
<b>IHC</b>	
<b>Reference</b>	Reference(s): Saitoh T, Moriwaki J, Koike J, Takagi A, Miwa T, Shiokawa K, Katoh M. Molecular cloning and characterization of FRAT2, encoding a positive regulator of the WNT signaling pathway. Biochem Biophys Res Commun. 2001 Mar 2;281(3):815-20..PMID: 11237732->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**