



Goat anti-CCKAR Antibody

Item Number	dAP-0798
Target Molecule	Principle Name: CCKAR; Official Symbol: CCKAR; All Names and Symbols: CCKAR; cholecystokinin A receptor ; HGNC:1570; CCK-A; CCK1-R; CCKRA; cholecystokinin type-A receptor; cholecystokinin-1 receptor; Accession Number (s): NP_000721.1; Human Gene ID(s): 886; Non-Human GeneID(s): 12425 (mouse) 24889 (rat)
Immunogen	FEASQKSAKERKPST, is from internal region
Applications	Pep ELISA, IHC, ICC, IF Species Tested: Human, Rat
Purification	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 4000.
Western Blot	Western Blot: Preliminary experiments in Human Pancreas lysates gave no specific signal but low background (at antibody concentration up to 1µg/ml).
IHC	Immunohistochemistry: In paraffin embedded Human Pancreas, shows cytoplasmic staining of some cells in the islets of Langerhans. Recommended concentration, 10µg/ml.
Reference	Reference(s): Beglinger C, Degen L, Matzinger D, D'Amato M, Drewe J. Loxiglumide, a CCK-A receptor antagonist, stimulates calorie intake and hunger feelings in humans. Am J Physiol Regul Integr Comp Physiol. 2001 Apr;280(4):R1149-54. .PMID: 11247838 ->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**