



## Goat anti-GLI3 Antibody

<b>Item Number</b>	dAP-0799
<b>Target Molecule</b>	Principle Name: GLI3; Official Symbol: GLI3; All Names and Symbols: GLI3; GLI-Kruppel family member GLI3 (Greig cephalopolysyndactyly syndrome) ; ACLS; GCPS; PAP-A; PAPA; PAPA1; PAPB; PHS; PPDIV ; DNA-binding protein; GLI-Kruppel family member GLI3; oncogene GLI3; zinc finger protein GLI3; Accession Number (s): NP_000159.2; Human Gene ID(s): 2737; Non-Human GeneID(s):
<b>Immunogen</b>	HNKRSKIKPDED, is from internal region
<b>Applications</b>	Pep ELISA Species Tested:
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 16000.
<b>Western Blot</b>	Western Blot: Preliminary experiments in Human Brain, Lung, Placenta and Testes lysates gave no specific signal but low background (at antibody concentration up to 1µg/ml). We would appreciate any feedback from people in the field - have any results been
<b>IHC</b>	
<b>Reference</b>	Reference(s): Haycraft CJ, Banizs B, Aydin-Son Y, Zhang Q, Michaud EJ, Yoder BK. Gli2 and gli3 localize to cilia and require the intraflagellar transport protein polaris for processing and function. PLoS Genet. 2005 Oct;1(4):e53. Epub 2005 Oct 28. .PMID: 16254602 ->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**