

DATA SHEET

## Goat anti-Ide (mouse) Antibody

Item Number	dAP-1128
Target Molecule	Principle Name: Ide (mouse); Official Symbol: Ide; All Names and Symbols: Ide; insulin-degrading enzyme; 1300012G03Rik; 4833415K22Rik; AA675336; AI507533; INSULYSIN; insulinase; insulysin; Accession Number (s): NP_112419.2; Human Gene ID(s): ; Non-Human GeneID(s): 15925 (mouse) 25700 (rat)
Immunogen	QQYNYDRDNIE, is from internal region
Applications	Pep ELISA, WB
Purification	Species Tested: Mouse, Rat Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Supplied As	lyophilized powder of 50ug or 100ug IgG; Reconsititute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
Peptide ELISA	Peptide ELISA: antibody detection limit dilution 1 to 16000.
Western Blot	Western Blot: Approx 110kDa band observed in Mouse Brain and Rat Brain lysates (calculated MW of 118kDa according to NP_112419.2). Recommended concentration: 1-3µg/ml. An additional band of unknown identity was also consistently observed at 30kDa. This
IHC	

Reference

Reference(s): Zhao Z, Xiang Z, Haroutunian V, Buxbaum JD, Stetka B, Pasinetti GM. Insulin degrading enzyme activity selectively decreases in the hippocampal formation of cases at high risk to develop Alzheimer's disease. Neurobiol Aging. 2007 Jun;28(6):824-30. Epub 2006 Jun 12..PMID: 16769157 ->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for Research Use Only