

## Goat anti-Lipocalin 2 / NGAL (mouse) Antibody

<b>Item Number</b>	dAP-1325
<b>Target Molecule</b>	Principle Name: Lipocalin 2 / NGAL (mouse); Official Symbol: Lcn2; All Names and Symbols: Lcn2; lipocalin 2; NGAL; RP23-161B9.11; 24p3; AW212229; NGAL; Accession Number (s): NP_032517.1; Human Gene ID(s): ; Non-Human GeneID(s): 16819 (mouse)
<b>Immunogen</b>	HRYPQVQSYNVQ , is from internal region
<b>Applications</b>	Pep ELISA, WB Species Tested: Mouse
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 8000.
<b>Western Blot</b>	Western Blot: Approx 23kDa band observed in Mouse Kidney lysates (calculated MW of 22.6kDa according to NP_005555.2). Recommended concentration: 1-3µg/ml.
<b>IHC</b>	
<b>Reference</b>	Reference(s): Venkatesha S, Hanai J, Seth P, Karumanchi SA, Sukhatme VP. Lipocalin 2 antagonizes the proangiogenic action of ras in transformed cells. Mol Cancer Res. 2006 Nov;4(11):821-9..PMID: 17114340 ->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**