

Goat anti-X-arrestin / arrestin 3 Antibody

| | |
|------------------------|--|
| Item Number | dAP-1771 |
| Target Molecule | Principle Name: X-arrestin / arrestin 3; Official Symbol: ARR3; All Names and Symbols: ARR3; arrestin 3, retinal (X-arrestin); RP13-26D14.6; ARRX; Arrestin C; Accession Number (s): NP_004303.2; Human Gene ID(s): 407; Non-Human GeneID(s): |
| Immunogen | RKGEEESQKAVE, is from C Terminus |
| Applications | Pep ELISA, IHC Species Tested: Human |
| Purification | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide. |
| Supplied As | lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing. |
| Peptide ELISA | Peptide ELISA: antibody detection limit dilution 1 to 1000. |
| Western Blot | |
| IHC | Immunohistochemistry: In paraffin embedded Human Retina shows staining of the outer segments of rod receptor cells. Recommended concentration, 2-4µg/ml. |

Reference Reference(s): Krishnamurthy H, Galet C, Ascoli M. The association of arrestin-3 with the follitropin receptor depends on receptor activation and phosphorylation. Mol Cell Endocrinol. 2003 Jun 30;204(1-2):127-40. PMID: 12850288 ->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**