

## Goat anti-ARHGDIB Antibody

<b>Item Number</b>	dAP-2178
<b>Target Molecule</b>	Principle Name: ARHGDIB; Official Symbol: ARHGDIB; All Names and Symbols: ARHGDIB; Rho GDP dissociation inhibitor (GDI) beta; D4; GDIA2; GDID4; LYGDI; Ly-GDI; RAP1GN1; RhoGDI2; Rho GDI 2; Rho GDP-dissociation inhibitor 2; Accession Number (s): NP_001166.3; Human Gene ID(s): 397; Non-Human GeneID(s):
<b>Immunogen</b>	TEKAPEPHVE, is from N Terminus
<b>Applications</b>	Pep ELISA, WB Species Tested: Human
<b>Purification</b>	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
<b>Supplied As</b>	lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.
<b>Peptide ELISA</b>	Peptide ELISA: antibody detection limit dilution 1 to 4000.
<b>Western Blot</b>	Western Blot: Approx 26kDa band observed in lysates of cell line Jurkat (calculated MW of 23.0kDa according to NP_001166.3). Recommended concentration: 1-3µg/ml.
<b>IHC</b>	

<b>Reference</b>	Reference(s): Guey LT, García-Closas M, Murta-Nascimento C, Lloreta J, Palencia L, Kogevinas M, Rothman N, Vellalta G, Calle ML, Marenne G, Tardón A, Carrato A, García-Closas R, Serra C, Silverman DT, Chanock S, Real FX, Malats N; EPICURO/Spanish Bladder Cancer Study i Genetic susceptibility to distinct
------------------	--

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**