



## **Goat anti-STAT5A Antibody**

Item Number dAP-2611

Target Molecule Principle Name: STAT5A; Official Symbol: STAT5A; All Names and Symbols: STAT5A; signal transducer

and activator of transcription 5A; MGF; STAT5; Accession Number (s): NP\_003143.2; Human Gene ID(s):

6776; Non-Human GeneID(s): 20850 (mouse) 24918 (rat)

Immunogen DSRLSPPAGLFTSAR, is from C Terminus

This antibody is not expected to cross-react with STAT5B.

Applications Pep ELISA, WB

Species Tested: Human

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

using the immunizing peptide.

Supplied As lyophilized powder of 50ug or 100ug lgG; Reconsititute lgG with 100ul or 200ul sterile DI Water and final

product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

Peptide ELISA Peptide ELISA: antibody detection limit dilution 1 to 128000.

Western Blot Western Blot: Approx 100kDa band observed in lysates of cell lines K562, Molt4 and NIH3T3 (calculated

MW of 90.6kDa according to NP 003143.2). Recommended concentration: 0.3-1µg/ml.

IHC

Reference Reference(s): Okutani Y, Kitanaka A, Tanaka T, Kamano H, Ohnishi H, Kubota Y, Ishida T,

Takahara J. Src directly tyrosine-phosphorylates STAT5 on its activation site and is involved in erythropoi-

etin-induced signaling pathway. Oncogene. 2001 Oct 4;20(45):6643-50..PMID: 11641791->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for Research Use Only