

## Goat anti-FSCN1 Antibody

|                        |   |
|------------------------|---|
| <b>Item Number</b>     | dAP-2950  |
| <b>Target Molecule</b> | Principle Name: FSCN1; Official Symbol: FSCN1 ; All Names and Symbols: FSCN1 ; fascin homolog 1, actin-bundling protein (Strongylocentrotus purpuratus) ; FLJ38511 ; SNL ; p55; Singed, drosophila, homolog-like ; actin bundling protein ; fascin 1 ; singed ; singed-like (fascin homolog, sea urchin); Accession Number (s): NP_003079.1; Human Gene ID(s): 6624; Non-Human GeneID(s): |
| <b>Immunogen</b>       | RDTKKCAFRTH, is from internal region  |
| <b>Applications</b>    | Pep ELISA, WB<br>Species Tested: Human  |
| <b>Purification</b>    | Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.   |
| <b>Supplied As</b>     | lyophilized powder of 50ug or 100ug IgG; Reconstitute IgG with 100ul or 200ul sterile DI Water and final product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.<br>Aliquot and store at -20°C. Minimize freezing and thawing.  |
| <b>Peptide ELISA</b>   | Peptide ELISA: antibody detection limit dilution 1 to 4000.   |
| <b>Western Blot</b>    | Western Blot: Approx 50kDa band observed in Human Skeletal Muscle lysates (calculated MW of 54.5kDa according to NP_003079.1). Recommended concentration: 1-3µg/ml.   |
| <b>IHC</b>             |   |
| <b>Reference</b>       | Reference(s): Vignjevic D, Kojima S, Aratyn Y, Danciu O, Svitkina T, Borisy GG. Role of fascin in filopodial protrusion. J Biol. 2006 Sep 11;174(6):863-75..PMID: 16966425 ->   |

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for **Research Use Only**