



## Goat anti-thyroid peroxidase, Biotinylated Antibody

Item Number dAP-3407

Target Molecule Principle Name: thyroid peroxidase, Biotinylated; Official Symbol: TPO; All Names and Symbols: TPO;

thyroid peroxidase; MSA; TDH2A; TPX; thyroid microsomal antigen; thyroperoxidase; Accession Number (s): NP\_000538.3; NP\_783650.1; NP\_783652.1; NP\_783653.1; Human Gene ID(s): 7173; Non-Human

GeneID(s):

Immunogen ELRCGKHQAVGTSPQ., is from C Terminus

This antibody is expected to recognize all reported isoforms (NP\_000538.3; NP\_783650.1; NP\_783652.1;

NP\_783653.1). Reported variants represent identical protein: NP\_000538.3, NP\_001193673.1 Reported

Applications Pep ELISA, WB, IHC

Species Tested: Human

Purification Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography

using the immunizing peptide.

Supplied As lyophilized powder of 50ug or 100ug lgG; Reconsititute lgG with 100ul or 200ul sterile DI Water and final

product will be formulated as 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum

albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

Peptide ELISA: antibody detection limit dilution 1 to 32000.

Western Blot: Approx 110kDa band observed in Human Thyroid Gland lysates (calculated MW of 103kDa

according to NP 000538.3). See non-biotinylated parental product's datasheet for further QC data. Recom-

mended concentration: 1-3µg/ml.

IHC

Reference Reference(s): Chen CR, Hamidi S, Braley-Mullen H, Nagayama Y, Bresee C, Aliesky HA, Rapoport B,

McLachlan SM. Antibodies to thyroid peroxidase arise spontaneously with age in NOD.H-2h4 mice and appear after thyroglobulin antibodies. Endocrinology 2010 Sep 151 (9): 4583-93. PMID: 20573721->

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for Research Use Only